

Submission

Inquiry into Sleep Health Awareness in Australia

The Impact of Artificial Light at Night on Sleep Health



October 2018

Summary

- Significant improvement in sleep health could be made by increasing public awareness that excessive artificial lighting at night, particularly in the blue end of the visual spectrum, is associated with poor sleep patterns and multiple medical conditions.
- Excessive exposure to artificial light at night is associated with other diseases including obesity, heart disease, diabetes, breast and prostate cancer
- Entities responsible for providing artificial lighting at night have a responsibility to limit such exposure on the public.
- Businesses that employ night shift workers, should be required to utilize lighting that minimizes the harmful effects of artificial lighting at night.
- Residential institutions such as nursing homes and hospitals should be required to utilize lighting that minimizes the harmful effects of artificial lighting at night.
- An added bonus of limiting artificial light at night is impact on other species, such as insects, migrating birds, night foragers and turtles.

Sleep wake cycles in animals and humans and some plants is controlled by melatonin, a hormone secreted by the brain. Melatonin levels are highest at night and suppressed during the daytime by exposure to blue rich light. In addition to its control of circadian rhythm, melatonin is an anti oxidant and plays a role in the immune system and suppression of tumour growth(1).

Exposure to artificial light, particularly in the blue area of the visual spectrum suppresses melatonin production, and, in turn causes sleep disturbance(2).

The light emitting diode (LED) lighting is becoming universal for electronic , indoor and outdoor lighting. LED lighting is very blue rich compared to more traditional lighting as it is essentially a blue LED with a phosphor coating to give the appearance of white light. Unless LED lighting is specifically filtered to minimize its blue light content it has more effect on Melatonin production and therefore sleep disturbance than more traditional lighting. Common sources of blue rich lights are, laptops, televisions, computer monitors, cool white house lights and cool white outdoor lighting, particularly street lighting (3-5).

The blue content of LEDS is often expressed as a kelvin rating. Where possible artificial lights at night should be rated no higher than 3000K. Ideally they should also have specific blue filters at the frequency that suppresses melatonin. The use of computer programs that decrease blue content on computers, phones and televisions after sunset, is recommended.

Solutions

Public awareness

While many factors contribute to sleep disturbance the raising of awareness of the above issue could be a relatively simple and cheap contribution to improving the nation's sleep health.

Outdoor lighting

While there is the psychological assumption that increased outdoor lighting is beneficial, there is no peer reviewed independent evidence that increasing outdoor lighting levels decreases crime rates, motor vehicle accidents or personal safety. Invariably articles that do support increased lighting suffer from poor scientific design, lack of peer review and/ or funded by lighting manufacturers.

The quality evidence shows that decreasing outdoor and street lighting does not lead to increases in motor vehicle accidents, crime rates or personal assaults(6).

There is a need to have providers of outdoor and public lighting contribute to minimization of exposure to artificial light at night. This was strongly recommended by the report of the American Medical Association(7).

Present standards often describe minimal lighting standards, but none describe maximum lighting standards. There is also a lack of standards requiring outdoor lighting to be sufficiently shielded so as not intrude on sleeping locations.

Occupational lighting

Businesses that employ night shift workers, should be required to utilize lighting that minimizes the harmful effects of artificial lighting at night.

Residential Institutions

Residential institutions such as nursing homes and hospitals, should be required to utilize lighting that minimizes the harmful effects of artificial lighting at night(8).

Since the introduction of the pulse oximeter to non invasively measure oxygen levels inpatients in 1986, there is no longer the imperative to have inpatients kept in a daytime lighting environment 24 hours a day, to detect cyanosed (blue skin) patients with low oxygen levels.

Staff in such institutions are particularly at risk from the damaging effects of suppressed melatonin secretion(9, 10).

Other conditions that would benefit from decreasing exposure to excessive artificial lighting at night.

1. Breast Cancer

Excessive artificial light is a recognized carcinogen(1, 9-15).

2. Prostate Cancer(2, 11, 14).
3. Obesity(2, 16).
4. Heart Disease(2, 16).
5. Diabetes(2, 16).
6. Depression(8)
7. Macular degeneration (17).
8. Impact on other species(18, 19).
9. Light pollution(20)

[REDACTED]

[REDACTED]

Appendix

1. Reiter RJ, Tan DX, Galano A. Melatonin: exceeding expectations. *Physiology*. 2014;29(5):325-33.
2. Cho Y, Ryu S-H, Lee BR, Kim KH, Lee E, Choi J. Effects of artificial light at night on human health: A literature review of observational and experimental studies applied to exposure assessment. *Chronobiology international*. 2015;32(9):1294-310.
3. Adams RJ, Appleton SL, Taylor AW, Gill TK, Lang C, McEvoy RD, et al. Sleep health of Australian adults in 2016: results of the 2016 Sleep Health Foundation national survey. *Sleep health*. 2017;3(1):35-42.
4. Emet M, Ozcan H, Ozel L, Yayla M, Halici Z, Hacimuftuoglu A. A review of melatonin, its receptors and drugs. *The Eurasian journal of medicine*. 2016;48(2):135.
5. Hall Ga. *Textbook of Medical Physiology*. 12th ed. Philadelphia US: Saunders Elsevier; 2011 2011.
6. Steinbach R, Perkins C, Tompson L, Johnson S, Armstrong B, Green J, et al. The effect of reduced street lighting on road casualties and crime in England and Wales: controlled interrupted time series analysis. *J Epidemiol Community Health*. 2015;69(11):1118-24.
7. Stevens R, Motta M. REPORT OF THE COUNCIL ON SCIENCE AND PUBLIC HEALTH. METHODS. 2018;23:24.
8. Obayashi K, Saeki K, Iwamoto J, Ikada Y, Kurumatani N. Exposure to light at night and risk of depression in the elderly. *Journal of affective disorders*. 2013;151(1):331-6.
9. Kloog I, Portnov BA, Rennert HS, Haim A. Does the modern urbanized sleeping habitat pose a breast cancer risk? *Chronobiology international*. 2011;28(1):76-80.
10. Kloog I, Stevens RG, Haim A, Portnov BA. Nighttime light level co-distributes with breast cancer incidence worldwide. *Cancer Causes & Control*. 2010;21(12):2059-68.
11. Garcia-Saenz A, Sanchez De Miguel A, Espinosa A, Valentin A, Aragonés N, Llorca J, et al. Evaluating the association between artificial light-at-night exposure and breast and prostate cancer risk in Spain (MCC-Spain study). 2018.
12. Greene MW. Circadian rhythms and tumor growth. *Cancer letters*. 2012;318(2):115-23.
13. Hahn RA. Profound bilateral blindness and the incidence of breast cancer. *Epidemiology*. 1991:208-10.
14. Pukkala E, Ojamo M, Rudanko S-L, Stevens RG, Verkasalo PK. Does incidence of breast cancer and prostate cancer decrease with increasing degree of visual impairment. *Cancer causes & control*. 2006;17(4):573-6.
15. Verkasalo P, Pukkala E, Stevens R, Ojamo M, Rudanko S-L. Inverse association between breast cancer incidence and degree of visual impairment in Finland. *British journal of cancer*. 1999;80(9):1459.

16. Falchi F, Cinzano P, Elvidge CD, Keith DM, Haim A. Limiting the impact of light pollution on human health, environment and stellar visibility. *Journal of environmental management*. 2011;92(10):2714-22.
17. Algvere PV, Marshall J, Seregard S. Age - related maculopathy and the impact of blue light hazard. *Acta Ophthalmologica Scandinavica*. 2006;84(1):4-15.
18. Tuxbury SM, Salmon M. Competitive interactions between artificial lighting and natural cues during seafinding by hatchling marine turtles. *Biological Conservation*. 2005;121(2):311-6.
19. Navara KJ, Nelson RJ. The dark side of light at night: physiological, epidemiological, and ecological consequences. *Journal of pineal research*. 2007;43(3):215-24.
20. Cinzano P, Falchi F, Elvidge CD. The first world atlas of the artificial night sky brightness. *Monthly Notices of the Royal Astronomical Society*. 2001;328(3):689-707.