

29 March 2016

Senator Alex Gallacher
Chair, Senate Foreign Affairs, Defence and Trade References Committee
Department of the Senate
PO Box 6100
Parliament House
CANBERRA ACT 2600

By email to: fadt.sen@aph.gov.au

Dear Senator Gallacher

Re: Australian Defence Force's resistance to interrogation training

The Royal Australian and New Zealand College of Psychiatrists (RANZCP) welcomes the opportunity to respond to the Senate Foreign Affairs, Defence and Trade Reference Committee's *Inquiry into the Australian Defence Force's (ADF) resistance to interrogation (RTI) training* (the Inquiry).

The RANZCP's response focuses on the Inquiry's term of reference (d): the ongoing mental health and wellbeing of personnel who have participated in RTI training.

Military service places distinctive mental health demands on ADF personnel. As noted in the *ADF Mental Health Prevalence and Wellbeing Study* (2010, p. 20):

Mental health and wellbeing in a military environment is unique. The military is an occupation where personnel are selected, trained and prepared to face adverse, stressful and potentially traumatising situations. To meet these demands, an approach that focuses on strengthening resilience and enabling recovery is essential.

At the same time, mental health issues affect many ADF personnel. For instance, 2010 research showed that one in five ADF members met the criteria for a mental disorder in the previous 12 months and the most common mental disorders in the ADF were anxiety disorders – in particular, post-traumatic stress disorder (PTSD) and affective (mood) disorders such as depressive episodes. The prevalence of suicidal ideation and making a suicide plan was also significantly higher among ADF personnel compared to the rest of the Australian community (ADF, 2010). This research suggests that 'despite the fact that the ADF is a selected and trained population that generally has better access to health care, this population bears a burden of psychiatric morbidity related to the nature of their work' (Van Hooff et al., 2014, p. 5).

Given the unique demands of military service, the RANZCP considers that it is essential that there are strategies to promote and support the mental health and wellbeing of current and former ADF personnel as well as services that provide them with quality mental health care. This includes for current and former ADF personnel, who have undertaken RTI training.

However, there is currently limited information available on the impact of RTI training on current and former ADF personnel and, therefore, it is an important area that requires further research.

The RANZCP also notes that stigma of mental illness is a major barrier for current and former ADF personnel seeking care for assistance with mental health problems. For instance, stigma attached to mental illness among military veterans may lead to denials and delays in care seeking, resulting in unnecessary suffering, self-medication, substance abuse and self-isolation (Sher and Braquehais, 2014).

Another related issue of concern is particular perception among ADF personnel that asking for help with mental health issues may impact detrimentally on their military careers. This is illustrated by the *ADF Mental Health Prevalence and Wellbeing Study*, which interviewed up to 49% of ADF members and reported that within a 12 month period: 27.6% were concerned that reporting a mental disorder might result in being treated differently, 26.9% feared their career might be harmed and 36.9% stated the main barrier to them seeking help was their concern that doing so would reduce the likelihood of them being deployed (ADF, 2010).

Therefore, the RANZCP welcomes the 11 June 2015 announcement that – following Phoenix Australia's review of ADF mental health screening processes - the ADF will be introducing new mental health screening tools later this year. These include providing an anonymous online self-screening mental health tool for serving members to access plus developing a mental health screening process, which will be used during routine medical appointments.

While these are welcome initiatives, the RANZCP remains concerned that the issue of stigma means that ADF personnel may not utilise these resources or, if they do, will understate any issues with their mental health. We consider systematic screening to be particularly important in identifying mental health issues, including those that might potentially arise due to ADF personnel's experiences with RTI. We, therefore, believe that it is crucial that the ADF implement an annual mental health survey during the time of ADF members' service and, after their service, every two to three years.

If you would like to discuss any of the issues raised in the submission, please contact

Yours sincerely

Professor Malcolm Hopwood
President

Ref: 0147o

References

Australian Defence Force (2010) Mental Health of the Australian Defence Force – 2010 ADF Mental Health Prevalence and Wellbeing Study Report. Available from:
www.defence.gov.au/Health/DMH/MentalHealthReformProgram.asp#MHRP.

Richardson L, Frueh, B C and Acierno, R (2010) Prevalence estimates of combat-related post-traumatic stress disorder: critical review. *Australian and New Zealand Journal of Psychiatry*. 44:4-19.

Sher, L and Braquehais, M D (2013) Suicidal behaviour in military veterans and health care professionals. *Australian and New Zealand Journal of Psychiatry* 48(6): 589.

Solomon, Z and Mikulincer, M (2006) Trajectories of PTSD: a 20 year longitudinal study. *American Journal of Psychiatry* 163: 659-666.

Van Hooff, M et al. (2014) The Australian Defence Force Mental Health Prevalence and Wellbeing Study: design and methods. *European Journal of Psychotraumatology*. 5: 10.3402/ejpt.v5.23950.