

## Appendix 1: Early Intervention in Psychosis – Selected Bibliography

- Access Economics, *Cost effectiveness of early intervention for psychosis*. 2008 , Access Economics: Canberra.
- Access Economics, *The economic impact of youth mental illness and the cost effectiveness of early intervention*. 2009, Access Economics: Canberra.
- Alvarez-Jimenez, M., et al., *Preventing the Second Episode: A Systematic Review and Meta-analysis of Psychosocial and Pharmacological Trials in First-Episode psychosis*. Schizophr Bull, 2009.
- Amminger, G.P., et al., *Long-chain omega-3 fatty acids for indicated prevention of psychotic disorders: a randomized, placebo-controlled trial*. Arch Gen Psychiatry, 2010. **67**(2): p. 146-154.
- Bechdolf, A., et al., *Randomized controlled multicentre trial of cognitive behavioural therapy in the early initial prodromal state: effects on social adjustment post treatment*. Early Interven Psychiatry, 2007. **1**(1): p. 71-78.
- Bertelsen, M., et al., *Five-year follow-up of a randomized multicenter trial of intensive early intervention vs standard treatment for patients with a first episode of psychotic illness: the OPUS trial*. Arch Gen Psychiatry, 2008. **65**(7): p. 762-771.
- Bertolote J and P.D. McGorry. *Early intervention and recovery for young people with early psychosis: Consensus statement*. Br J Psychiatry 2010. **187** (suppl.48): s. 116-119
- Birchwood, M., P. Todd, and C. Jackson, *Early intervention in psychosis. The critical period hypothesis*. Br J Psychiatry Suppl, 1998. **172**(33): p. 53-59.
- Bird V., et al, *Early intervention services, cognitive behavioural therapy and family intervention in early psychosis: systemic review*. Br J Psychiatry 2010. **197** (5): p 350-356
- Bowers, M., *The onset of psychosis--a diary account*. Psychiatry, 1965. **28**(4): p. 346-358.
- Cannon, T.D., et al., *Prediction of psychosis in youth at high clinical risk: a multisite longitudinal study in North America*. Arch Gen Psychiatry, 2008. **65**(1): p. 28-37.
- Caton, C.L., et al., *Stability of early-phase primary psychotic disorders with concurrent substance use and substance-induced psychosis*. Br J Psychiatry, 2007. **190**: p. 105-111.
- Chapman, L.J. and J.P. Chapman, *The search for symptoms predictive of schizophrenia*. Schizophr Bull, 1987. **13**(3): p. 497-503.
- Chen, E.Y., et al., *Maintenance treatment with quetiapine versus discontinuation after one year of treatment in patients with remitted first episode psychosis: randomised controlled trial*. BMJ, 2010. **341**: p. c4024.
- Cornblatt, B., T. Lencz, and M. Obuchowski, *The schizophrenia prodrome: treatment and high-risk perspectives*. Schizophr Res, 2002. **54**(1-2): p. 177-186.
- Cornblatt, B.A., et al., *Can antidepressants be used to treat the schizophrenia prodrome? Results of a prospective, naturalistic treatment study of adolescents*. J Clin Psychiatry, 2007. **68**(4): p. 546-557.
- Craig, T.K., et al., *The Lambeth Early Onset (LEO) Team: randomised controlled trial of the effectiveness of specialised care for early psychosis*. BMJ, 2004. **329**(7474): p. 1067.
- Cullberg, J., et al., *Treatment costs and clinical outcome for first episode schizophrenia patients: a 3-year follow-up of the Swedish "Parachute Project" and two comparison groups*. Acta Psychiatr Scand, 2006. **114**(4): p. 274-281.
- Dodgson, G., et al., *Early intervention in psychosis service and psychiatric admissions*. The Psychiatrist, 2008. **32**: p. 413-416.

- Early Psychosis Guidelines Writing Group, *Australian clinical guidelines for early psychosis*. 2nd edition ed. 2010, Melbourne: Orygen Youth Health.
- Edwards, J. and P.D. McGorry, *Implementing early intervention in psychosis: a guide to establishing early psychosis services*. 2002, London: Martin Dunitz.
- Edwards, J., et al., *Randomized controlled trial of clozapine and CBT for first-episode psychosis with enduring positive symptoms: a pilot study*. Schizophr Res Treat, 2011, in press.
- Emsley, R.A., *Risperidone in the treatment of first-episode psychotic patients: a double-blind multicenter study*. Risperidone Working Group. Schizophr Bull, 1999. **25**(4): p. 721-729.
- Fowler, D., et al., *Can targeted early intervention improve functional recovery in psychosis? A historical control evaluation of the effectiveness of different models of early intervention service provision in Norfolk 1998–2007*. Early Interven Psychiatry, 2009. **3**: p. 282-288.
- Friis, S., *Early specialised treatment for first-episode psychosis: does it make a difference?* Br J Psychiatry, 2010. **196**(5): p. 339-340.
- Fusar-Poli, P., L. Valmaggia, and P. McGuire, *Can antidepressants prevent psychosis?* Lancet, 2007. **370**(9601): p. 1746-1748.
- Gaebel, W., et al., *Relapse prevention in first-episode schizophrenia-maintenance vs intermittent drug treatment with prodrome-based early intervention: results of a randomized controlled trial within the german research network on schizophrenia*. J Clin Psychiatry, 2010..
- Gafoor, R., et al., *Effect of early intervention on 5-year outcome in non-affective psychosis*. Br J Psychiatry, 2010. **196**(5): p. 372-376.
- Garety, P.A., et al., *Specialised care for early psychosis: symptoms, social functioning and patient satisfaction: randomised controlled trial*. Br J Psychiatry, 2006. **188**: p. 37-45.
- Gleeson, J., T.K. Larsen, and P. McGorry, *Psychological treatment in pre- and early psychosis*. J Am Acad Psychoanal Dyn Psychiatry, 2003. **31**(1): p. 229-245.
- Gleeson, J.F., et al., *Early warning signs of relapse following a first episode of psychosis*. Schizophr Res, 2005. **80**(1): p. 107-111.
- Goldberg, K., et al., *Impact of a specialized early intervention service for psychotic disorders on patient characteristics, service use, and hospital costs in a defined catchment area*. Can J Psychiatry, 2006. **51**(14): p. 895-903.
- Grawe, R.W., et al., *Two years of continued early treatment for recent-onset schizophrenia: a randomised controlled study*. Acta Psychiatr Scand, 2006. **114**(5): p. 328-336.
- Haddock, G., et al., *A pilot study evaluating the effectiveness of individual inpatient cognitive-behavioural therapy in early psychosis*. Soc Psychiatry Psychiatr Epidemiol, 1999. **34**(5): p. 254-258.
- Hafner, H. and K. Maurer, *Early detection of schizophrenia: current evidence and future perspectives*. World Psychiatry, 2006. **5**(3): p. 130-138.
- Harrigan, S.M., P.D. McGorry, and H. Krstev, *Does treatment delay in first-episode psychosis really matter?* Psychol Med, 2003. **33**(1): p. 97-110.
- Harris, M.G., et al., *Impact of a specialized early psychosis programme on suicide. A retrospective cohort study*. Early Interven Psychiatry, 2008. **2**: p. 11-21.

Harris, M.G., et al., *The relationship between duration of untreated psychosis and outcome: an eight-year prospective study*. Schizophr Res, 2005. **79**(1): p. 85-93.

Heinrichs, D.W. and W.T. Carpenter, Jr., *Prospective study of prodromal symptoms in schizophrenic relapse*. Am J Psychiatry, 1985. **142**(3): p. 371-373.

International Early Psychosis Association Writing Group, *International clinical practice guidelines for early psychosis*. Br J Psychiatry Suppl, 2005. **48**: p. s120-124.

Jackson, C. and M. Birchwood, *Early intervention in psychosis: opportunities for secondary prevention*. Br J Clin Psychol, 1996. **35** (4): p. 487-502.

Jackson, H.J., et al., *Acute-phase and 1-year follow-up results of a randomized controlled trial of CBT versus Befriending for first-episode psychosis: the ACE project*. Psychol Med, 2008. **38**(5): p. 725-735.

Jeppesen, P., et al., *Integrated treatment of first-episode psychosis: effect of treatment on family burden: OPUS trial*. Br J Psychiatry Suppl, 2005. **48**: p. s85-90.

Joa, I., et al., *The key to reducing duration of untreated first psychosis: information campaigns*. Schizophr Bull, 2008. **34**(3): p. 466-472.

Johannessen, J.O., et al., *Early detection strategies for untreated first-episode psychosis*. Schizophr Res, 2001. **51**(1): p. 39-46.

Johannessen, J.O., et al., *Pathways to care for first-episode psychosis in an early detection healthcare sector: part of the Scandinavian TIPS study*. Br J Psychiatry Suppl, 2005. **48**: p. s24-28.

Jorgensen, P., et al., *Early detection and assertive community treatment of young psychotics: the Opus Study Rationale and design of the trial*. Soc Psychiatry Psychiatr Epidemiol, 2000. **35**(7): p. 283-287.

Kahn, R.S., et al., *Effectiveness of antipsychotic drugs in first-episode schizophrenia and schizophreniform disorder: an open randomised clinical trial*. Lancet, 2008. **371**(9618): p. 1085-1097.

Killackey, E., H.J. Jackson, and P.D. McGorry, *Vocational intervention in first-episode psychosis: individual placement and support v. treatment as usual*. Br J Psychiatry, 2008. **193**(2): p. 114-120.

Killackey, E.J., et al., *Exciting career opportunity beckons! Early intervention and vocational rehabilitation in first-episode psychosis: employing cautious optimism*. Aust N Z J Psychiatry, 2006. **40**(11-12): p. 951-962.

Kuipers, E., et al., *An RCT of early intervention in psychosis: Croydon Outreach and Assertive Support Team (COAST)*. Soc Psychiatry Psychiatr Epidemiol, 2004. **39**(5): p. 358-363.

Lambert, M., et al., *Pharmacotherapy of first-episode psychosis*. Expert Opin Pharmacother, 2003. **4**(5): p. 717-750.

Larsen, T.K., et al., *Early detection of first-episode psychosis: the effect on 1-year outcome*. Schizophr Bull, 2006. **32**(4): p. 758-764.

Larsen, T.K., et al., *Early detection of psychosis: positive effects on 5-year outcome*. Psychol Med: p. 1-9.

Lecomte, T., et al., *Group cognitive behavior therapy or social skills training for individuals with a recent onset of psychosis? Results of a randomized controlled trial*. J Nerv Ment Dis, 2008. **196**(12): p. 866-875.

Lewis, S., et al., *Randomised controlled trial of cognitive-behavioural therapy in early schizophrenia: acute-phase outcomes*. Br J Psychiatry Suppl, 2002. **43**: p. s91-97.

Marshall, M., et al., *Association between duration of untreated psychosis and outcome in cohorts of first-episode patients: a systematic review*. Arch Gen Psychiatry, 2005. **62**(9): p. 975-983.

McCrone, P., et al., *Cost-effectiveness of an early intervention service for people with psychosis*. Br J Psychiatry, 2010. **196**(5): p. 377-382.

McCrone, P., M. Knapp, and S. Dhanasiri, *Economic impact of services for first-episode psychosis: a decision model approach*. Early Interv Psychiatry, 2009. **3**: p. 266-273.

McGlashan, T.H., *Early detection and intervention in psychosis: an ethical paradigm shift*. Br J Psychiatry Suppl, 2005. **48**: p. s113-115.

McGlashan, T.H., et al., *Randomized, double-blind trial of olanzapine versus placebo in patients prodromally symptomatic for psychosis*. Am J Psychiatry, 2006. **163**(5): p. 790-799.

McGlashan, T.H., et al., *The PRIME North America randomized double-blind clinical trial of olanzapine versus placebo in patients at risk of being prodromally symptomatic for psychosis. I. Study rationale and design*. Schizophr Res, 2003. **61**(1): p. 7-18.

McGlashan, T.H., *Rationale and parameters for medication-free research in psychosis*. Schizophr Bull, 2006. **32**(2): p. 300-302.

McGorry, P., M. Nordentoft, and E. Simonsen, *Introduction to "Early psychosis: a bridge to the future"*. Br J Psychiatry Suppl, 2005. **48**: p. s1-3.

McGorry, P.D. and A.R. Yung, *Early intervention in psychosis: an overdue reform*. Aust N Z J Psychiatry, 2003. **37**(4): p. 393-398.

McGorry, P.D., A.R. Yung, and L.J. Phillips, *The "close-in" or ultra high-risk model: a safe and effective strategy for research and clinical intervention in prepsychotic mental disorder*. Schizophr Bull, 2003. **29**(4): p. 771-790.

McGorry, P.D., E. Killackey, and A. Yung, *Early intervention in psychosis: concepts, evidence and future directions*. World Psychiatry, 2008. **7**(3): p. 148-156.

McGorry, P.D., E. Killackey, and A.R. Yung, *Early intervention in psychotic disorders: detection and treatment of the first episode and the critical early stages*. Med J Aust, 2007. **187**(7 Suppl): p. S8-10.

McGorry, P.D., et al., *Clinical staging of psychiatric disorders: a heuristic framework for choosing earlier, safer and more effective interventions*. Aust N Z J Psychiatry, 2006. **40**(8): p. 616-622.

McGorry, P.D., et al., *Clinical staging: a heuristic and practical strategy for new research and better health and social outcomes for psychotic and related mood disorders*. Can J Psychiatry, 2010. **55**(8): p. 486-497.

McGorry, P.D., et al., *Clinical staging: a heuristic model for psychiatry and youth mental health*. Med J Aust, 2007. **187**(7 Suppl): p. s40-42.

McGorry, P.D., et al., *EPPIC: an evolving system of early detection and optimal management*. Schizophr Bull, 1996. **22**(2): p. 305-326.

McGorry, P.D., et al., *High risk/prodrome/early intervention*, in *The Year in Schizophrenia*, G. Thaker and W.T. Carpenter, Editors. 2007, Clinical Publishing: London. p. 45-57.

McGorry, P.D., et al., *High risk/prodrome/early intervention*, in *The Year in Schizophrenia*, G. Thaker and W.T. Carpenter, Editors. 2008, Clinical Publishing: London. p. 55-70.

McGorry, P.D., et al., *Randomized controlled trial of interventions designed to reduce the risk of progression to first-episode psychosis in a clinical sample with subthreshold symptoms*. Arch Gen Psychiatry, 2002. **59**(10): p. 921-928.

- McGorry, P.D., *Issues for DSM-V: clinical staging: a heuristic pathway to valid nosology and safer, more effective treatment in psychiatry*. Am J Psychiatry, 2007. **164**(6): p. 859-860.
- Melle, I., et al., *Early detection of the first episode of schizophrenia and suicidal behavior*. Am J Psychiatry, 2006. **163**(5): p. 800-804.
- Melle, I., et al., *Reducing the duration of untreated first-episode psychosis: effects on clinical presentation*. Arch Gen Psychiatry, 2004. **61**(2): p. 143-150.
- Mihalopoulos, C., et al., *Is early intervention in psychosis cost-effective over the long term?* Schizophr Bull, 2009. **35**(5): p. 909-918.
- Mihalopoulos, C., P.D. McGorry, and R.C. Carter, *Is phase-specific, community-oriented treatment of early psychosis an economically viable method of improving outcome?* Acta Psychiatr Scand, 1999. **100**(1): p. 47-55.
- Moller, P. and R. Husby, *The initial prodrome in schizophrenia: searching for naturalistic core dimensions of experience and behavior*. Schizophr Bull, 2000. **26**(1): p. 217-232.
- Morrison, A.P., et al., *Cognitive therapy for the prevention of psychosis in people at ultra-high risk: randomised controlled trial*. Br J Psychiatry, 2004. **185**: p. 291-297.
- Morrison, A.P., et al., *Three-year follow-up of a randomized controlled trial of cognitive therapy for the prevention of psychosis in people at ultrahigh risk*. Schizophr Bull, 2007. **33**(3): p. 682-687.
- Mrazek, P.J. and R.J. Haggerty, *Reducing risks for mental disorders: frontiers for preventive intervention research*. 1994, Washington: National Academy Press.
- NHS Confederation, *Briefing*, 2011 Issue 219, NHS Confederation: London, UK
- Nielsen, O. and M. Large, *Rates of homicide during the first episode of psychosis and after treatment: a systematic review and meta-analysis*. Schizophr Bull, 2010. **36**(4): p. 702-712.
- Nordentoft, M., et al., *Transition rates from schizotypal disorder to psychotic disorder for first-contact patients included in the OPUS trial. A randomized clinical trial of integrated treatment and standard treatment*. Schizophr Res, 2006. **83**(1): p. 29-40.
- Nordentoft, M., L. Bjarking, and R. Hemmingsen, *Psychiatric emergency outreach: a report on the first 2 years of functioning in Copenhagen*. Nord J Psychiatry, 2002. **56**(6): p. 399-405.
- Norman et al 2011 *Symptomatic and functional outcomes for a 5 year early intervention program for psychoses* Schizophr Res, 2011. **129** p111-115
- Patton, G.C., S.E. Hetrick, and P. McGorry, *Service responses for youth onset mental disorders*. Curr Opin Psychiatry, 2007. **20**(4): p. 319-324.
- Perez-Iglesias, R., et al., *Effect of antipsychotics on peptides involved in energy balance in drug-naive psychotic patients after 1 year of treatment*. J Clin Psychopharmacol, 2008. **28**(3): p. 289-295.
- Perkins, D.O., et al., *Relationship between duration of untreated psychosis and outcome in first-episode schizophrenia: a critical review and meta-analysis*. Am J Psychiatry, 2005. **162**(10): p. 1785-1804.
- Petersen, L., et al., *A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness*. BMJ, 2005. **331**(7517): p. 602.

Petersen, L., et al., *Improving 1-year outcome in first-episode psychosis: OPUS trial*. Br J Psychiatry Suppl, 2005. **48**: p. s98-103.

Phillips, L.J., et al., *Medium term follow-up of a randomized controlled trial of interventions for young people at ultra high risk of psychosis*. Schizophr Res, 2007. **96**(1-3): p. 25-33.

Phillips, L.J., et al., *Randomized controlled trial of interventions for young people at ultra-high risk of psychosis: study design and baseline characteristics*. Aust N Z J Psychiatry, 2009. **43**(9): p. 818-829.

Poulton, R., et al., *Children's self-reported psychotic symptoms and adult schizophreniform disorder: a 15-year longitudinal study*. Arch Gen Psychiatry, 2000. **57**(11): p. 1053-1058.

Rahm, C. and J. Cullberg, *Diagnostic stability over 3 years in a total group of first-episode psychosis patients*. Nord J Psychiatry, 2007. **61**(3): p. 189-193.

Rinaldi, M., et al., *What are the benefits of evidence-based supported employment for patients with first-episode psychosis?* Psychiatr Bull, 2004. **28**(8): p. 281-284.

Robinson, D.G., et al., *Pharmacological treatments for first-episode schizophrenia*. Schizophr Bull, 2005. **31**(3): p. 705-722.

Robinson, D.G., et al., *Predictors of treatment response from a first episode of schizophrenia or schizoaffective disorder*. Am J Psychiatry, 1999. **156**(4): p. 544-549.

Robinson, J., et al., *Suicide attempt in first-episode psychosis: a 7.4 year follow-up study*. Schizophr Res, 2010. **116**(1): p. 1-8.

Ruhrmann, S., F. Schultze-Lutter, and J. Klosterkötter, *Early detection and intervention in the initial prodromal phase of schizophrenia*. Pharmacopsychiatry, 2003. **36** (Suppl 3): p. s162-167.

Salvatore, P., et al., *McLean-Harvard International First-Episode Project: two-year stability of DSM-IV diagnoses in 500 first-episode psychotic disorder patients*. J Clin Psychiatry, 2009. **70**(4): p. 458-466.

Sanger, T.M., et al., *Olanzapine versus haloperidol treatment in first-episode psychosis*. Am J Psychiatry, 1999. **156**(1): p. 79-87.

Schimmelmann, B.G., et al., *Diagnostic stability 18 months after treatment initiation for first-episode psychosis*. J Clin Psychiatry, 2005. **66**(10): p. 1239-1246.

Singh, S.P., *Early intervention in psychosis*. Br J Psychiatry, 2010. **196**(5): p. 343-345.

Tarrier, N., et al., *Cognitive-behavioural therapy in first-episode and early schizophrenia. 18-month follow-up of a randomised controlled trial*. Br J Psychiatry, 2004. **184**: p. 231-239.

Thorup, A., et al., *Integrated treatment ameliorates negative symptoms in first episode psychosis--results from the Danish OPUS trial*. Schizophr Res, 2005. **79**(1): p. 95-105.

Tien, A.Y. and J.C. Anthony, *Epidemiological analysis of alcohol and drug use as risk factors for psychotic experiences*. J Nerv Ment Dis, 1990. **178**(8): p. 473-480.

Valmaggia, L.R., et al., *Economic impact of early intervention in people at high risk of psychosis*. Psychol Med, 2009. **39**(10): p. 1617-1626.

Verdoux, H. and A. Cougnard, *Schizophrenia: who is at risk? Who is a case?* Int Clin Psychopharmacol, 2006. **21** (Suppl 2): p. s17-19.

Victorian Department of Health Mental Health Branch, *Youth Early Psychosis Status Report*. 2007, Victorian Department of Health: Melbourne.

Whitty, P., et al., *Diagnostic stability four years after a first episode of psychosis*. Psychiatr Serv, 2005. **56**(9): p. 1084-1088.

- Woods, S.W., et al., *Validity of the prodromal risk syndrome for first psychosis: findings from the North American Prodrome Longitudinal Study*. Schizophr Bull, 2009. **35**(5): p. 894-908.
- Wunderink, L., et al., *Guided discontinuation versus maintenance treatment in remitted first-episode psychosis: relapse rates and functional outcome*. J Clin Psychiatry, 2007. **68**(5): p. 654-661.
- Yung, A.R. and P.D. McGorry, *Prediction of psychosis: setting the stage*. Br J Psychiatry Suppl, 2007. **51**: p. s1-8.
- Yung, A.R. and P.D. McGorry, *The initial prodrome in psychosis: descriptive and qualitative aspects*. Aust N Z J Psychiatry, 1996. **30**(5): p. 587-599.
- Yung, A.R. and P.D. McGorry, *The prodromal phase of first-episode psychosis: past and current conceptualizations*. Schizophr Bull, 1996. **22**(2): p. 353-370.
- Yung, A.R., et al., *Declining transition rate in ultra high risk (prodromal) services: dilution or reduction of risk?* Schizophr Bull, 2007. **33**(3): p. 673-681.
- Yung, A.R., et al., *Monitoring and care of young people at incipient risk of psychosis*. Schizophr Bull, 1996. **22**(2): p. 283-303.
- Yung, A.R., et al., *Prediction of psychosis. A step towards indicated prevention of schizophrenia*. Br J Psychiatry Suppl, 1998. **172**(33): p. 14-20.
- Yung, A.R., et al., *Psychosis prediction: 12-month follow up of a high-risk ("prodromal") group*. Schizophr Res, 2003. **60**(1): p. 21-32.
- Yung, A.R., et al., *Risk factors for psychosis in an ultra high-risk group: psychopathology and clinical features*. Schizophr Res, 2004. **67**(2-3): p. 131-142.